



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Objective functional testing in patients with lumbar degenerative disc disease

Joswig, Holger ; Stienen, M N ; Smoll, N R ; Gautschi, O P

DOI: <https://doi.org/10.1177/2192568217716153>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-133209>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Joswig, Holger; Stienen, M N; Smoll, N R; Gautschi, O P (2017). Objective functional testing in patients with lumbar degenerative disc disease. *Global Spine Journal*, 7(4):384.

DOI: <https://doi.org/10.1177/2192568217716153>

Objective Functional Testing in Patients With Lumbar Degenerative Disc Disease

Holger Joswig, MD¹, Martin N. Stienen, MD²,
Nicolas R. Smoll, MD³, and Oliver P. Gautschi, MD²

Global Spine Journal
2017, Vol. 7(4) 384
© The Author(s) 2017
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/2192568217716153
journals.sagepub.com/home/gsj



Dear Editor:

We were delighted to read the results of the prospective study by Mobbs et al¹ on “Physical Activity Measured With Accelerometer and Self-Rated Disability in Lumbar Spine Surgery” in a recent issue of the *Global Spine Journal*. Using a commercially available accelerometer and patient-reported outcome measures (PROMs), such as the Visual Analog Scale (VAS), Oswestry Disability Score (ODI), and Short-Form-12 (SF-12) questionnaire, the authors evaluated improvement of objective and subjective function in $n = 30$ patients following lumbar spine surgery including laminectomy, discectomy, and fusion. Lack of correlation of the investigated PROMs with changes in the physical activity parameters (steps, distance, and calories) and the exceptional ability of the accelerometer as an objective tool to determine objective function has encouraged the authors to recommend the inclusion of objective physical activity measurements. Despite the small cohort, the authors should be congratulated for their convincing results, which hopefully will stimulate more clinical research in this field.

The authors appraised to be “the first to prospectively investigate objective physical activity measurements after lumbar spine surgery and test whether these measurements correlate with subjective functional scores.” However, it is worth mentioning that we called for the need for an objective outcome measurement in spine surgery as early as in October 2014,² and thus introduced the Timed Up and Go (TUG) test as a measurement tool of objective functional impairment (OFI) in the context of lumbar degenerative disc disease (DDD) and its surgical treatment.³ In short, the TUG test measures the time in seconds to stand up from a chair, walk 3 meters, turn around, walk 3 meters back, and sit down again. These physical functions are essential to perform the majority of activities of daily living. Using a stopwatch, a health care professional can easily determine the presence and degree of OFI in patients, stratified into mild, moderate, and severe OFI. Alternatively, the results can be presented as standardized age- and sex-adjusted t scores.³ For the evaluation of a patient in the postoperative course, a

reduction of the TUG test time of at least 3.4 seconds translated into a clinically meaningful improvement.⁴ Recently, a TUG smartphone application has been developed to measure the raw TUG test times, OFI, and t scores in patients with lumbar DDD.

We are in full agreement with the statement made by Mobbs et al¹ that objective assessment tools should find their way into clinical research and practice. In our experience, the objective assessment adds a new dimension to the armamentarium of clinical tests for a comprehensive patient evaluation, but does not replace well-established PROMs that capture important—subjective—aspects.

References

1. Mobbs RJ, Phan K, Maharaj M, Rao PJ. Physical activity measured with accelerometer and self-rated disability in lumbar spine surgery: a prospective study. *Global Spine J*. 2016;6:459-464.
2. Gautschi OP, Corniola MV, Schaller K, Smoll NR, Stienen MN. The need for an objective outcome measurement in spine surgery—the timed-up-and-go test. *Spine J*. 2014;14:2521-2522.
3. Gautschi OP, Smoll NR, Corniola MV, et al. Validity and reliability of a measurement of objective functional impairment in lumbar degenerative disc disease: the Timed Up and Go (TUG) test. *Neurosurgery*. 2016;79:270-278.
4. Gautschi OP, Stienen MN, Corniola MV, et al. Assessment of the minimum clinically important difference in the Timed Up and Go Test after surgery for lumbar degenerative disc disease. *Neurosurgery*. 2017;80:380-385.

¹ University Hospital, London, Ontario, Canada

² University Hospital Geneva, Geneva, Switzerland

³ John Hunter Hospital, Newcastle, Australia

Corresponding Author:

Martin N. Stienen, MD, Department of Neurosurgery, University Hospital Zurich, Frauenklinikstrasse 10, 8091 Zürich, Switzerland.
Email: mnstienen@gmail.com



This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 License (<http://www.creativecommons.org/licenses/by-nc-nd/4.0/>) which permits non-commercial use, reproduction and distribution of the work as published without adaptation or alteration, without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).